

cleanslate Knoxville

Disinfectant + Sanitizer + Virucide

- Kills SARS-CoV-2, which causes COVID-19² on hard non-porous surfaces in just 30 seconds!
- Kills 99.9%¹ of bacteria and viruses* on hard, non-porous surfaces.
- Formulated for third sink¹

¹SARS-Related Coronavirus 2 ²When used according to disinfection directions.

¹ Formulated for 3rd sink food contact surface sanitization.



Escanee el código QR para obtener declaraciones de precaución e instrucciones de almacenamiento /eliminación en español.

KEEP OUT OF REACH OF CHILDREN DANGER PELIGRO

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. **IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. **IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. **IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

This product for use in: restaurants, fast food operations, bars, food storage areas, food establishments, coffee shops, bagel stores, food processing plants and meat/poultry processing plants. This product is for use as a sanitizer on dishes, glassware and utensils at 200-400 ppm active quaternary without potable water rinse. This product is for use as a sanitizer in public eating establishments and on food processing equipment at 200-400 ppm active quaternary without potable water rinse. Sanitization of public eating establishment food contact surfaces is regulated under 40CFR180.940. This product is for use as a hard, non-porous surface hospital disinfectant at 547 ppm active quaternary and as a general disinfectant at 469 ppm active quaternary.

ACTIVE INGREDIENTS

Alkyl (60% C14, 30% C16, 5% C12, 5% C18) dimethyl benzyl ammonium chlorides.....5.0%
Alkyl (68% C12, 32% C14) dimethyl ethylbenzyl ammonium chlorides.....5.0%
INERT INGREDIENTS.....90.0%

TOTAL.....100.0%

EPA Reg. No. 1839-86-84179 EPA Est. No. 84179-NJ-1

Sold by: Cleanslate Group LLC 1515 East Linden Ave, Linden NJ 07036

net contents:
1 Gallon (128 fl oz)

USE: For visibly soiled areas, a pre-cleaning step is required. Prepare a fresh solution for each use. To disinfect inanimate, hard non-porous surfaces apply use-solution with mop, cloth, sponge, low pressure coarse sprayer or hand pump trigger sprayer so as to wet all surfaces thoroughly. Allow to remain visibly wet for 10 minutes, then remove excess liquid. Surfaces which may contact food must be rinsed thoroughly with potable water after use. For low pressure coarse sprayer and hand pump trigger sprayer applications, spray 6-8 inches from the surface; rub with brush, sponge or cloth. Do not breathe spray. **NOTE:** For spray applications, cover or remove all food products. **GENERAL DISINFECTION -** Add 3 fluid ounces of this product per 5 gallons of water for disinfection against *Staphylococcus aureus*, *Salmonella enterica*, *Listeria monocytogenes*, and *Yersinia enterocolitica*. **HOSPITAL DISINFECTION -** Add 3.5 fluid ounces of this product per 5 gallons of water for disinfection against *Pseudomonas aeruginosa* and Vancomycin intermediate resistant *Staphylococcus aureus* (VISA), as well as the organisms listed under the GENERAL DISINFECTION section. **To disinfect food processing premises:** floors, walls, and storage areas, add 3 fluid ounces of this product per 5 gallons of water. For visibly soiled areas, a pre-cleaning step is required. Apply solution with a mop, cloth, sponge or hand pump trigger sprayer so as to wet all surfaces thoroughly. Allow to remain visibly wet for 10 minutes, then remove excess liquid. Before using this product, food products and packaging materials must be removed from the area or carefully protected. After use, all surfaces in the area must be thoroughly rinsed with potable water. **To disinfect food service establishment or restaurant food contact surfaces:** floors, walls, storage areas, countertops, outside of appliances, and tables, add 3 fluid ounces of this product per 5 gallons of water. For visibly soiled areas, a pre-cleaning step is required. Apply solution with a cloth, sponge or hand pump trigger sprayer so as to wet all surfaces thoroughly. Allow the surface to remain visibly wet for 10 minutes, then remove excess liquid and rinse the surface with potable water. Do not use on utensils, dishes, glasses or cookware. ***VIRUCIDAL ACTIVITY -** This product, when used on environmental, inanimate, hard non-porous surfaces at 3.5 fluid ounces per 5 gallons (or equivalent dilution) of water exhibits effective virucidal activity against Influenza A2 Virus (representative of the common flu virus), Herpes Simplex Virus Type 1 (causative agent of fever blisters), Adenovirus Type 5 (causative agent of upper respiratory infections), Vaccinia Virus (representative of the pox virus group), Avian Influenza A Virus (causative agent of an acute avian lower respiratory tract infection), HIV-1 (Human Immunodeficiency Virus Type 1), Newcastle Disease Virus, Laryngotracheitis Virus, Hepatitis B Virus (HBV), Hepatitis C Virus (HCV), Bovine Viral Diarrhea Virus (BVD), Human Coronavirus (ATCC VR-740, Strain 229E), Porcine Respiratory & Reproductive Syndrome Virus (PRRSV) (Strain NVSL) and SARS associated Coronavirus. **SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV AND HCV OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS. PERSONAL PROTECTION:** When handling items soiled with blood or body fluids use disposable latex gloves, gowns, masks, and eye coverings. **CLEANING PROCEDURES:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product. **CONTACT TIME:** Allow surface to remain visibly wet for 10 minutes. **DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal. **SARS-CoV-2 ACTIVITY:** At 2 fluid ounces per gallon of water (or equivalent dilution) this product is an effective virucide against the following viruses on hard non-porous environmental surfaces: SARS-CoV-2 after 30 seconds. **FOOD CONTACT SURFACE SANITIZATION DIRECTIONS Sanitization:** Recommended for use in restaurants, food processing plants, and bars. When used as directed this product is an effective sanitizer at an active quaternary concentration of 200-400 ppm when diluted in water up to 650 ppm hardness (CaCO3) against *Escherichia coli* (ATCC 11229), *Escherichia coli* O157:H7 (ATCC 43895), *Staphylococcus aureus* (ATCC 6538), *Yersinia enterocolitica* (ATCC 23715), *Listeria monocytogenes* (ATCC 35152), *Salmonella typhi* (ATCC 6539), *Shigella sonnei* (ATCC 11060), *Vibrio cholera* (ATCC 14035), Methicillin resistant *Staphylococcus aureus* (MRSA) (ATCC 33592), Vancomycin resistant *Enterococcus faecalis* (VRE) (ATCC 51299), and *Campylobacter jejuni* (ATCC 29428). This product is an effective sanitizer against *Klebsiella pneumoniae* (ATCC 4352) at a 200 ppm active quaternary concentration when diluted in water up to 500 ppm hardness (CaCO3). At an active quaternary concentration of 300 ppm, it is an effective sanitizer against *Klebsiella pneumoniae* (ATCC 4352) when diluted in water up to 650 ppm hardness (CaCO3). Remove all visible food particles and soil from areas, which are to be sanitized, with a good detergent, pre-flush, pre-soak or pre-scrub treatment. Rinse with a potable water rinse. Prepare a 200-400 ppm active quaternary solution by adding 1-2 fluid ounces of this product to 4 gallons of water, 2.5-5 fluid ounces to 10 gallons of water or 5-10 fluid ounces to 20 gallons of water. Prepare a fresh solution daily or more frequently as soil is apparent. **To sanitize pre-cleaned mobile food contact utensils** (such as drinking glasses, dishes, eating utensils), immerse in a 200-400 ppm active quaternary solution for at least 60 seconds making sure to immerse completely. Remove items, drain the use-solution from the surface and air dry. Do not rinse. **To sanitize pre-cleaned food contact surfaces too large for immersion** (such as counters, tables, finished wood or plastic cutting boards), apply a 200-400 ppm active quaternary solution with a cloth, sponge, low pressure coarse sprayer or hand pump trigger sprayer making sure that the surface remains completely visibly wet for at least 60 seconds. Drain the use-solution from the surface and let air dry. Do not rinse. **Sanitization of food processing equipment, utensils, and other food contact articles regulated under 40CFR180.940:** 1. Scrape, flush, or pre-soak articles (whether mobile or stationary) to remove visible food particles and soil. 2. Thoroughly wash articles with a compatible detergent or cleaner. 3. Rinse articles thoroughly with potable water. 4. Sanitize by immersing articles with a use-solution of 1-2 fluid ounces of this product per 4 gallons of water (200-400 ppm active quaternary) for at least 60 seconds. Articles too large for immersion should be thoroughly wetted by rinsing, spraying or swabbing. 5. Remove immersed items from solution to drain and then let air dry. Non-immersed items must be allowed to air dry.

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL. PESTICIDE STORAGE: Store in a dry place no lower in temperature than 50°F or higher than 120°F. Store in a tightly closed container in an area inaccessible to children. **PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER DISPOSAL:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container promptly after emptying. Triple rinse as follows: Fill container full with water and recap. Agitate vigorously. Follow Pesticide Disposal instruction for rinsate disposal. Drain for 10 seconds after the flow begins to drip. Repeat procedure two more times. Offer for recycling or reconditioning, if available. If not available, puncture and dispose in a sanitary landfill. **SPILL OR LEAK PROCEDURES:** Small spills may be mopped up, flushed away with water or absorbed on some absorbent material and incinerated. Large spills should be contained, the material then moved into containers and disposed of by approved methods for hazardous wastes.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS.

DANGER. KEEP OUT OF REACH OF CHILDREN. CORROSIVE. Causes irreversible eye damage and skin burns. Do not get in eyes, on skin or on clothing. Wear goggles or face shield, rubber gloves, and protective clothing. Harmful if swallowed. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling.